

# HACKENSACK DIGESTIVE DISEASE ASSOCIATES, P.A.

BOARD CERTIFIED IN GASTROENTEROLOGY & LIVER DISEASE \*

**PLEASE FOLLOW THESE INSTRUCTIONS CAREFULLY.** They are extremely important to the success and accuracy of your examination, as well as for your safety. If you have any questions, please contact our office at 201-488-3003.

Your colonoscopy is scheduled at...

Hackensack Endoscopy Center  
(Excelsior II Building)  
**170 Prospect Avenue**  
Suite 10  
Hackensack, NJ 07601

Endoscopy Center of Bergen County  
**80 Eisenhower Drive**  
Paramus, NJ 07652

Hackensack Medical Center  
Medical Plaza Building  
**20 Prospect Avenue**  
Suite 813  
Hackensack, NJ 07601

Your appointment is scheduled for \_\_\_\_\_ at \_\_\_\_\_.

**FOLLOW THE INSTRUCTIONS ON THE ENCLOSED SHEET ON HOW TO TAKE YOUR BOWEL PREPARATION EXACTLY AS DIRECTED.**

Please do not consume any raw fruits or vegetables for 3 days prior to your exam.

**The day before you colonoscopy you cannot eat any solid food and you must follow a clear liquid diet.** Clear liquids include Gatorade (not red), clear broth, apple juice, Jell-O (not red), black coffee (no milk or cream), tea, water, soda, popsicles and Italian ice (not red).

**You cannot have anything to eat or drink after midnight the night before your colonoscopy.** You can take medications, such as blood pressure medication, on the day of your procedure with a sip of water.

Please do not take aspirin, Ecotrin or anti-inflammatory medication, such as Motrin (ibuprofen) or Aleve, for one week prior to your colonoscopy, unless otherwise instructed by your physician.

If you take **Coumadin** (warfarin), **Plavix** (clopidogrel), **Pradaxa**, **Xarelto**, **Eliquis** or **Effient**, please contact our office if you have not already received specific instructions regarding taking these medications and your planned colonoscopy.

If you take insulin or other diabetes medications, please contact our office or the doctor who prescribed these medications for specific instructions.

You will be receiving sedation for your colonoscopy and **YOU CANNOT DRIVE YOURSELF HOME.** You must arrange for someone to drive you home after your colonoscopy.

The day before your colonoscopy at 4:00 PM, begin drinking the Nulytely/Colyte. Drink 8 ounces every 10-15 minutes. This process could take up to four hours to complete. You may continue drinking clear liquids after you have completed the Nulytely/Colyte until midnight. Do not drink anything after midnight.